

OXYPEST

COPPER OXYCHLORIDE 50% WP

FUNGICIDE

- **OXYPEST** BELONGS TO INORGANIC GROUP.
- **OXYPEST** IS A FOLIAR FUNGICIDE WITH CONTACT AND PREVENTIVE ACTION.
- **OXYPEST** BELONGS TO INORGANIC GROUP.
- **OXYPEST** IS HIGHLY SOLUBLE IN WATER AND PRODUCES UNIFORM DISTRIBUTION ON PLANT

AVAILABLE PACK SIZES:

500G, 1KG

ANTIDOTE:

DIMERCAPROL IS GIVEN IN THE DOSE OF 3 MG/KG EVERY FOUR HOURS FOR FIRST FOUR DAYS FULLY ATROPINISED.



RECOMMENDATION FOR USE:

CROP	DISEASE	DOSAGE
		KG/ACRE
BANANA	LEAF SPOT, FRUIT ROT	1
BETEL	LEAF SPOT, FRUIT ROT	1
CHILLI	FRUIT ROT, LEAF SPOT	1
CITRUS	CANKER, LEAF SPOT	1
CARDAMOM	CLUMP ROT	1.5-2
	LEAF SPOT	1
COFFEE	BLACK ROT, RUST	1.5-2
COCONUT	BUD ROT	1
POTATO	EARLY BLIGHT, LATE	1